

Magnesium Sulphate - Management of Hypertensive Disorders of Pregnancy



1. Purpose

Magnesium sulphate is the anticonvulsant of choice for pre-eclampsia prophylaxis and treatment.

This clinical guideline outlines the indications, contraindications, administration and monitoring of Magnesium sulphate use at the Women's.

Also refer to Procedure: [Observations – Birth Centre – Adult Escalation Criteria and Response Framework](#).

2. Definitions

Not applicable.

3. Responsibilities

Staff caring for the patient with pre- eclampsia, eclampsia should be aware about this guideline.

4. Guideline

4.1 Indications for use

- magnesium sulphate is used in pregnancy for women with pre-eclampsia for whom there is concern about the risk of eclampsia
- firstline management of an eclamptic seizure
- neuroprotection of preterm infants.

4.2 Presentation

10mL = 500mg/mL (50% solution) (*magnesium sulfate heptahydrate*) and;

50mL vials = 493 mg/mL (49.3% treat as ~ 50% solution).

4.3 Route of administration

IV infusion:

- magnesium sulphate is delivered by controlled infusion via a syringe pump.
- the infusion should be connected via a multi-flow adapter to a peripheral line of normal saline 0.9% and monitored by an infusion pump in order to reduce localized irritation and monitor fluid balance.
- The magnesium line should be labelled clearly and not used to inject other medicines.

Note: magnesium sulphate infusions should only be administered in Birth Centre, Theatre or Complex Care Unit.

4.4 Dose

Prophylaxis of pre-eclampsia seizure

Loading dose: using a 10mL vial of magnesium sulphate prepare 4gram (i.e. **8mL**) of magnesium sulphate 50% in a 10mL syringe, configure the pump to accept the 10mL syringe and set the pump to 32mL an hour for 15 minutes.

Maintenance rate: once the loading dose has been completed, using the 50mL vial of magnesium sulphate, prepare 50mL of magnesium sulphate 50% in a 50mL syringe, re-set the pump to accept 50mL syringe and set the pump to administer the maintenance rate of 1g/hr (2mL/hour) or as ordered, until at least 24 hours post birth/delivery.

NB: Ensure magnesium sulphate is administered concurrently via a Y-site with a compatible IV fluid.

4.5 Side effects:

- hypotension secondary to reductions in systemic vascular resistance
- facial flushing

Magnesium Sulphate - Management of Hypertensive Disorders of Pregnancy



the women's
the royal women's hospital

- visual disturbances
- flushing at injection site
- chest pain
- nasal stuffiness.

NOTE: caution should be taken with any additional intravenous fluid administration to manage hypotension because of the potential risk of pulmonary oedema.

The following may also occur:

- ECG changes
- circulatory collapse
- gastro-intestinal upset
- urinary retention
- magnesium toxicity
- tissue necrosis at the injection site.

4.6 Contraindications and precautions

- Magnesium sulphate should be administered with caution in women being treated with cardiac glycosides/digitalis.
- Concurrent use of magnesium sulphate and CNS depressants may result in an enhanced CNS depressant effect.
- If calcium gluconate is required to treat magnesium sulphate toxicity, it should be delivered with caution as it may precipitate heart block.
- As magnesium is excreted in urine, patients with impaired renal function or electrolyte imbalance should be managed cautiously and magnesium levels should be monitored closely because of the increased risk of magnesium toxicity.

4.7 Initial and subsequent management of eclamptic seizures

The priorities are to terminate the seizure and prevent maternal and fetal hypoxia.

The first line of treatment for a woman who is having an eclamptic seizure is supportive care which should follow the basic principles for maintaining the woman's airway, breathing and circulation.

Refer to: CPG: Eclampsia: Management.

- Call for help and initiate: Code Blue - Adult and Child (intranet-only).
- Do not leave the woman alone. Assess the situation in regards to safety of staff and patient.
- Aim to prevent maternal injury during the seizure.
- Position the woman into the left lateral position and administer oxygen by mask.
- Continue to assess AIRWAY, BREATHING, CIRCULATION
- Secure IV access.
- Check maternal pulse and blood pressure. Use pulse oximetry if available.

Magnesium sulphate is the most effective medicine for the treatment of seizures when administered to women with eclampsia.

Magnesium Sulphate - Management of Hypertensive Disorders of Pregnancy



AVOID POLYPHARMACY to treat seizures as this will increase the risk of respiratory arrest.

Note: If a maintenance infusion is in progress, cease the infusion.

Give LOADING DOSE:

- draw up 4g (8mL) of magnesium sulphate 50% in 10mL syringe
- administer intravenously over ten (10) minutes – slow IV push

Commence/re-commence maintenance infusion.

REMEMBER to check ALL pump limits when recommencing the infusion.

Note: ECG monitoring should be available and anaesthetists should be aware of the patient's medical condition.

Recurrent seizures should be treated with a further bolus dose of 4g (8mL) magnesium sulphate.

4.8 Monitoring

Magnesium level monitoring

The therapeutic level of magnesium: 1.7 to 3.5 mmol/L. Routine monitoring of magnesium levels is not required BUT as magnesium is excreted by the kidneys, monitoring of serum levels should be conducted in women with renal impairment (6 hourly or more frequently as indicated).

Request magnesium level and review management if:

- respiratory rate < 12 breaths/minute
- urine output < 100mLs in 4 hours
- loss of patellar reflexes
- further seizures occur.

Mg conc (mmol/L)	Effects
0.8 - 1.0	normal plasma level
1.7 - 3.5	therapeutic range
2.5 - 5.0	ECG changes (P-Q interval prolongation, widen QRS complex)
4.0 - 5.0	reduction in deep tendon reflexes
> 5.0	loss of deep tendon reflexes
> 7.5	sinoatrial and atrioventricular blockade. Respiratory paralysis and CNS depression
> 12	cardiac arrest

Note: If serum magnesium level is >3.5mmol/L, cease infusion and consult with obstetrician.

Clinical observations

During administration of the loading or bolus dose:

- 5 minutely blood pressure and pulse (x 4 readings)
- observe for the development of side effects
- check patellar reflexes after administration.

Magnesium Sulphate - Management of Hypertensive Disorders of Pregnancy



4.9 Clinical practice note: taking of maternal blood pressure

The woman should be resting at an angle of greater than 45 degrees with her feet supported.

The blood pressure cuff should be of the appropriate size and should be placed at the level of the heart.

Standard cuff for arms <33cm circumference. Large cuff (15 x 33cm bladder) for larger arms.

Inflate cuff to 20-30mmHg above palpated systolic pressure. Deflate slowly.

Read and record blood pressure to nearest 2mmHg.

Korotkoff phase 5 sound (sound disappearance) is the appropriate measurement of diastolic blood pressure.

Multiple levels should be taken to confirm the diagnosis of hypertension/ pre-eclampsia due to natural variation.

Automated blood pressure readings should be used with caution in pre-eclampsia. They can systematically underestimate, more particularly, the systolic blood pressure.

It is recommended that a calibrated manual sphygmomanometer be used initially as a reference unless it is known that the automatic machine has been validated for pregnancy.

During administration of the maintenance infusion:

- ½ hourly blood pressure, pulse, and respiratory rate (pre-treatment respiratory rate should be ≥ 16per minute). These may be undertaken hourly post-birth.
- 1 hourly patellar reflexes
- 1 hourly urine measures, 4 hourly testing of urinary protein
- 2 hourly temperature
- continuous electronic fetal monitoring from 26 weeks gestation until clinical review/discussion by medical staff. Between 24- 26 weeks gestation, individualised management with regard to fetal monitoring will be considered
- maintain strict fluid balance chart.

Record patellar reflexes as:

A = Absent

N = Normal

B = Brisk

Response to magnesium toxicity

The following clinical signs of magnesium toxicity must be reviewed by a consultant obstetrician/anaesthetist:

- urine output <100mL in 4 hours
- absent patellar reflexes
- respiratory depression.

The antidote for magnesium toxicity is: 10mL calcium gluconate (available as 2.2mmol of calcium in 10mL vial- formerly known as 10% solution) over 10 minutes by slow intravenous injection. The patient requires ECG monitoring during and after administration because of the potential for cardiac arrhythmias.

Resuscitation and ventilator support should be available during and after dose administration of both magnesium sulphate and calcium gluconate.

CEASE Magnesium infusion in the following emergencies:

respiratory arrest: call: Code Blue - Adult and Child (intranet-only)

Magnesium Sulphate - Management of Hypertensive Disorders of Pregnancy



cardiac arrest: call: Code Blue - Adult and Child (intranet-only)

Ceasing a magnesium infusion

Magnesium infusion to continue for a minimum of 24 hours post birth/delivery (continue hourly clinical observation for 4 hours following the discontinuation of magnesium infusion). Cease according to medical orders.

Postpartum magnesium levels may be adequately assessed clinically (reflexes, respiratory rate) unless there is renal impairment/oliguria when serum levels should be continued 6 hourly.

5. Evaluation, monitoring and reporting of compliance to this guideline

Compliance to this guideline will be monitored by review of incidents reported through VHIMS.

6. References

Refer to the learning package: [Magnesium Sulphate Infusions](#).

Women's Policies, Guidelines and Procedures:

- [Eclampsia Management](#)
- [Pre-Eclampsia: Management](#)

7. Legislation/Regulations related to this guideline

Not applicable

8. Appendices

Not applicable.

PGP Disclaimer Statement

The Royal Women's Hospital Clinical Guidelines present statements of 'Best Practice' based on thorough evaluation of evidence and are intended for health professionals only. For practitioners outside the Women's this material is made available in good faith as a resource for use by health professionals to draw on in developing their own protocols, guided by published medical evidence. In doing so, practitioners should themselves be familiar with the literature and make their own interpretations of it.

Whilst appreciable care has been taken in the preparation of clinical guidelines which appear on this web page, the Royal Women's Hospital provides these as a service only and does not warrant the accuracy of these guidelines. Any representation implied or expressed concerning the efficacy, appropriateness or suitability of any treatment or product is expressly negated

In view of the possibility of human error and / or advances in medical knowledge, the Royal Women's Hospital cannot and does not warrant that the information contained in the guidelines is in every respect accurate or complete. Accordingly, the Royal Women's Hospital will not be held responsible or liable for any errors or omissions that may be found in any of the information at this site.

You are encouraged to consult other sources in order to confirm the information contained in any of the guidelines and, in the event that medical treatment is required, to take professional, expert advice from a legally qualified and appropriately experienced medical practitioner.

NOTE: Care should be taken when printing any clinical guideline from this site. Updates to these guidelines will take place as necessary. It is therefore advised that regular visits to this site will be needed to access the most current version of these guidelines.

Magnesium Sulphate - Management of Hypertensive Disorders of Pregnancy

