

OVULATION INDUCTION PROGRAM AT THE WOMEN'S



the women's
the royal women's hospital
victoria australia

Ovulation Induction (OI) is a treatment option for couples who have been unable to conceive despite regular unprotected intercourse over a 6–12 month period.

Sometimes an imbalance in a woman's reproductive hormones leads to irregular or absent menstrual cycles. This can make getting pregnant difficult.

Women referred to the **Ovulation Induction Program** are prescribed medications to help address this imbalance. The medications stimulate the growth of ovarian follicles and assist ovulation.

While you are on the Program

Any medications you may need are available from the Women's Pharmacy on Level 1 and from pharmacies in the community. If you wish to purchase the medications from your local pharmacy you will need to check if they stock them.

You will be monitored to assess your response to the medications. Monitoring may involve;

- » one or more vaginal ultrasounds to assess ovarian follicle growth
- » using urinary Luteinizing Hormone (LH) detection kits to identify when ovulation occurs
- » blood tests to check hormone levels.

It is important to have regular intercourse during the 'fertile' time in your cycle. Intercourse with your partner in the days up to and including the day of ovulation will give you the best chance of pregnancy.

Starting your OI cycle

Please telephone the OI Nurse on the first day of your period (full flow of bleed). This is called Day One. The OI nurse will help you plan your cycle, i.e. when to start taking your medication and the dose required. The OI nurse will also book you in for your first ultrasound appointment.

About the medications and tests

Clomifene tablets

Clomifene tablets are usually started early in your cycle (between day three and five) and are taken for five days. This medication stimulates the release of naturally occurring FSH, which will cause ovarian follicles to grow.

Letrozole

Letrozole tablets are also started early in the cycle (between day three and five) and are taken for five days. Letrozole increases the production of FSH by suppressing the production of oestrogen. The production of FSH will cause ovarian follicles to grow.

FSH Injections – Gonal F® or Puregon®

FSH injections are a course of daily injections administered at the same time each day. Your first injections will be on day three of your cycle.

Clomiphene and FSH may cause your ovaries to grow more than one follicle which increases your risk of having a multiple pregnancy. Talk to your doctor or nurse about this risk, they will help you decide whether to continue or cancel your OI cycle. See also *Ovulation induction – multiple pregnancy risk*.

Vaginal ultrasound

Vaginal ultrasounds measure the thickness of the lining of the uterus (the endometrium) as well as the size and number of follicles on both ovaries. This scan can be done by one of the doctors or nurses at Reproductive Services (RSU).

You may need more than one vaginal scan before you are ready to be triggered for ovulation. The number of scans depends entirely on the time taken for your body to grow follicles. If you are on daily FSH injections, the dose of FSH may need to change depending on your scan results.

According to hospital policy you may be scanned by a male or female doctor/nurse. RSU will try and meet your request to be scanned by a female practitioner, but this cannot be guaranteed.

Urinary LH detection kits

After your first vaginal ultrasound you may be asked to start home urine LH tests. The test detects your body's LH surge, which occurs 24 hours before ovulation. Home urine testing kits can be purchased through the clinic. Your urine needs to be tested daily.

Trigger injection – Human Chorionic Gonadotrophin – Pregnyl® 5000IU

Trigger injections cause ovulation to occur 38 to 42 hours after administration. It is a one-off injection given in the same way as FSH injections. Some patients 'surge' spontaneously, (i.e. begin ovulating) so do not need this injection. Your OI nurse will advise you on what date it should be administered.

The trigger injection can be given in the clinic by a nurse or self-administered at home. The timing is not crucial so can be given at any time of the day. It is important to have regular intercourse once ovulation has been triggered.

Luteal Phase Support (LPS)

The luteal phase is the part of a woman's cycle that begins with ovulation and ends with a period. During the luteal phase the body produces more oestrogen and progesterone which work together to change the lining of the uterus and prepare it to accept the implantation of an embryo.

Your doctor may order additional medications to help keep the uterine lining thick and discourage an early period. This is referred to as Luteal Phase Support.

The medication routinely used for LPS is Pregnyl® 1500IU. Two injections are required four and eight days after ovulation.

If you have used Clomifene or Letrozole tablets earlier in the cycle it is unlikely that LPS will be ordered. Your nurse will advise you if any medication is necessary and on what dates to administer the LPS injections.

Progesterone test

In your first OI cycle, a blood test will be required approximately seven days after ovulation to measure your progesterone levels. Progesterone is an important hormone in the luteal phase, and the test gives your doctor information about whether your levels are sufficient. If the test shows that your progesterone levels are lower than expected your doctor may need to adjust your medications in future cycles.

Pregnancy test

If you do not have a period within 19 days of ovulation an appointment for a pregnancy blood test must be scheduled. Please phone Reproductive Services Reception who will book an appointment for you.

If the test is taken early in the morning your results are generally available the same day. If you prefer to have your blood test done locally, we can arrange a request form for a local pathology provider.

What if I get my period?

If you unfortunately have a full flow normal period after an OI cycle, please phone the OI nurse to notify us of your Day One, a new cycle of OI will be planned for you.

If you do bleed but the period is not normal for you please phone the OI nurses, we may organise a pregnancy test as your OI cycle may still have been successful.

Timing of intercourse

The ideal time for intercourse is before and up to the time of ovulation. Intercourse may commence any time after the first scan. Once a large follicle (>16mm) is identified you should have frequent intercourse (every couple of days).

DISCLAIMER The Royal Women's Hospital does not accept any liability to any person for the information or advice (or use of such information or advice) which is provided in this fact sheet or incorporated into it by reference. We provide this information on the understanding that all persons accessing it take responsibility for assessing its relevance and accuracy. Women are encouraged to discuss their health needs with a health practitioner. If you have concerns about your health, you should seek advice from your health care provider or if you require urgent care you should go to the nearest hospital Emergency Department. © The Royal Women's Hospital, June 2018

Review

A review appointment should be made with a doctor in Reproductive Services if you have had no success after three ovulatory cycles.

Common medication side effects

Clomifene tablets (Clomid and Serophene)

- » Headache
- » Depression
- » Hot flushes
- » Temporary visual disturbances like blurring and yellow discolouration

Letrozole

Hot flushes

- » Joint pain
- » Fatigue
- » Nausea

FSH (Gonal f® or Puregon®)

- » Abdominal distension and discomfort
- » Increased vaginal discharge
- » Breast tenderness
- » Enlarged, tender ovaries
- » Bruising or irritation at injection site
- » Irritability, tearfulness
- » Headaches
- » Nausea
- » Fluid retention

HCG (Pregnyl® 5000IU and 1500IU)

- » Nausea
- » Vomiting
- » Weight gain
- » Shortness of breath
- » Diarrhoea
- » Painful breasts
- » Bloating
- » Mild stomach pain

Contact information at the Women's

Reproductive Services

Level 2, Royal Women's Hospital

Reception: (03) 8345 3200

OI Nurse: (03) 8345 3230 - Monday to Friday 7.30am to 3.30pm

Email: oinurse@thewomens.org.au

Pharmacy Department

Level 1, Royal Women's Hospital

Tel: (03) 8345 3180